

Fritz-Haber-Institut der Max-Planck-Gesellschaft, Humboldt-Universität zu Berlin, Max-Delbrück-Centrum für Molekulare Medizin, Otto-von-Guericke-Universität Magdeburg, Physikalisch-Technische Bundesanstalt, Technische Universität Berlin, Universität Potsdam



Berlin Center for Studies of Complex Chemical Systems

Seminar

Complex Nonlinear Processes in Chemistry and Biology

Honorary Chairman: G. Ertl

Organizers: M. Bär, C. Beta, H. Engel, M. Falcke, M. J. B. Hauser, J. Kurths, A. S. Mikhailov,

P. Plath, L. Schimansky-Geier, and H. Stark

Friday, June 2, 2017, at 10:00 c.t.

Address: Universität Potsdam, Institut für Physik und Astronomie, Karl-Liebknecht-Str. 24/25, 14476 Potsdam-Golm, Haus 28, Room 2.28.0.108

Dr. Michael Stich

Aston University (Birmingham, UK)

Symmetry breaking in simple models of cooperative polymerization

Biological homochirality is the well-established preference of biomolecules for only one of two chiral isomers (or enantiomers), as for example observed in sugars (D-enantiomer) and amino acids (L-enantiomer). Understanding how this symmetry breaking may have emerged remains a challenge for origin of life research. After discussing some fundamental experimental and theoretical findings, we investigate a kinetic rate equation model of nucleated cooperative enantioselective polymerization in closed systems [1]. The microreversible scheme includes (i) solution-phase racemization of the monomers, (ii) linear homochiral chain growth by stepwise monomer attachment and (iii) annealing or fusion of chains. Mechanically induced breakage of the longest chains maintains the system out of equilibrium and drives a breakage-fusion recycling mechanism. Spontaneous mirror symmetry breaking can be achieved starting from small initial enantiomeric excesses due to the intrinsic statistical fluctuations. We put our findings into the context of simpler models, trying to understand the onset of symmetry breaking.

[1] C. Blanco, M. Stich, D. Hochberg, J. Phys. Chem. B **121**, 942 (2017).